#### (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization International Bureau



## 

## (43) International Publication Date 25 October 2001 (25.10.2001)

#### **PCT**

# (10) International Publication Number WO 01/78737 A1

- (51) International Patent Classification?: A61K 31/575, 31/167, A61P 11/06 // (A61K 31/575, 31:167)
- (21) International Application Number: PCT/GB01/01628
- (22) International Filing Date: 11 April 2001 (11.04.2001)
- (25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data: 0009584.4

18 April 2000 (18.04.2000) GE

- (71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): GAVIN, Brian, Charles [IE/IE]; GlaxoSmithKline, P.O. Box 700, Grange Road, Rathfarnfam, 16 Dublin (IE).
- (74) Agent: LEAROYD, Stephanie, Anne; GlaxoSmithKline, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.



#### MEDICAL COMBINATIONS COMPRISING FORMOTEROL AND BUDESONIDE

The present invention is concerned with combinations of (R,R)-formoterol and budesonide, particularly compositions containing a combination of (R.R)formoterol and budesonide and the use of such compositions in medicine. particularly in the prophylaxis and treatment of respiratory diseases.

Formoterol, i.e. 2'-hydroxy-5'-[(RS)-1-hydroxy-2{[(RS)-p-methoxy-\alphamethylphenethyl]amino}ethyl]formanilide, particularly its fumarate salt is a wellknown adrenoreceptor agonist which is now used clinically in the treatment of bronchial asthma and related disorders. Formoterol includes two asymmetric centres and in a particular form exists as the (R,R)- isomer. The (R,R) isomer of formoterol has been described previously, for example, in WO98/21175 and US5795564.

15

10

5

DE 2,323,215 and US 3,929,768 describe budesonide i.e.  $(11\beta,16\alpha)$ -16,17-[butylidenebis(oxy)]-11,21-dihydroxypregna-1,4-diene-3,20-dione, salts thereof and pharmaceutical formulations thereof. Budesonide is an antiinflammatory corticosteroid, which is now used clinically in the treatment of bronchial asthma and related disorders.

20

WO 93/11773 describes combinations of budesonide and formoterol but is silent as to the utility of (R,R)-formoterol.

25 Although (R,R)-formoterol fumarate and budesonide are effective therapies, there exists a clinical need for asthma therapies having potent and selective action and having an advantageous profile of action.

Therefore, according to the present invention there is provided a combination of 30 (R,R)-formoterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof.

10

15

20

25

30

It will be appreciated that the compounds of the combination may be administered simultaneously, either in the same or different pharmaceutical formulations or sequentially. If there is sequential administration, the delay in administering the second compound should not be such as to lose the beneficial therapeutic effect of the combination.

According to a further aspect of the present invention, there is provided a pharmaceutical formulation comprising (R,R)-formoterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients. According to a preferred aspect of the present invention, there is provided a pharmaceutical formulation comprising (R,R)-formoterol fumarate and budesonide, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients. In the most preferred aspect, the above pharmaceutical formulations are suitable for administration by inhalation.

It is to be understood that the present invention covers all combinations of particular and preferred aspects of the invention described herein.

By the term "physiologically functional derivative" is meant a chemical derivative of (R,R)-formoterol or budesonide having the same physiological function as the free compound, for example, by being convertible in the body thereto. According to the present invention, examples of physiologically functional derivatives include esters.

Suitable salts according to the invention include those formed with both organic and inorganic acids. Pharmaceutically acceptable acid addition salts include but are not limited to those formed from hydrochloric, hydrobromic, sulphuric, citric, tartaric, phosphoric, lactic, pyruvic, acetic, trifluoroacetic, succinic, oxalic, fumaric, maleic, oxaloacetic, methanesulphonic, ethanesulphonic, ptoluenesulphonic, benzenesulphonic, isethionic, and naphthalenecarboxylic, such as 1-hydroxy-2-naphthalenecarboxylic acids.

Pharmaceutically acceptable esters of (R,R)-formoterol or budesonide may have a hydroxyl group converted to a  $C_{1-6}$  alkyl, aryl, aryl  $C_{1-6}$  alkyl, or amino acid ester.

As mentioned above, both (R,R)-formoterol and budesonide and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have been described for use in the treatment of respiratory diseases. Therefore, formulations of (R,R)-formoterol and budesonide and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have use in the prophylaxis and treatment of clinical conditions for which a selective β<sub>2</sub>-adrenoreceptor agonist and/or an antiinflammatory corticosteroid is indicated. Such conditions include diseases associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary diseases (COPD) (e.g. chronic and wheezy bronchitis, emphysema), respiratory tract infection and upper respiratory tract disease.

20

25

30

35

Accordingly, the present invention provides a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which a selective β2-adrenoreceptor agonist and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a combination of (R,R)-formoterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof. The present invention further provides a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which a selective β<sub>2</sub>-adrenoreceptor agonist and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a pharmaceutical formulation comprising (R,R)-formoterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient. In a preferred aspect, there is provided such a method which comprises administration of a therapeutically effective amount of a pharmaceutical formulation comprising (R,R)-formoterol furnarate and budesonide, and a pharmaceutically acceptable carrier or excipient. In

4

particular, the present invention provides such methods for the prophylaxis or treatment of a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

5

10

15

20

35

In the alternative, there is provided a combination of (R,R)-formoterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, for use in therapy, particularly for use in the prophylaxis or treatment of a clinical condition for which a selective β<sub>2</sub>-adrenoreceptor agonist and/or antiinflammatory corticosteroid is indicated. In particular, there is provided a pharmaceutical formulation comprising (R,R)formoterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof (suitably, (R,R)-formoterol fumarate) and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient for use in therapy, particularly for use in the prophylaxis or treatment of a clinical condition for which a selective β2-adrenoreceptor agonist and/or antiinflammatory corticosteroid is indicated. In a preferred aspect, the invention is concerned with the prophylaxis or treatment of a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

The amount of (R,R)-formoterol and budesonide, or a pharmaceutically
acceptable salt, solvate or physiologically functional derivative thereof which is
required to achieve a therapeutic effect will, of course, vary with the particular
compound, the route of administration, the subject under treatment, and the
particular disorder or disease being treated. As a monotherapy, (R,R)-formoterol
fumarate is generally administered to adult humans by aerosol inhalation at a
dose of 12mcg or 24mcg twice daily. As a monotherapy, budesonide is
generally administered to adult humans by aerosol inhalation at a dose of from
200mcg to 1.6mg daily, taken as 2 divided doses:

While it is possible for the active ingredients of the combination to be administered as the raw chemical, it is preferable to present them as a

WO 01/78737

5

PCT/GB01/01628

pharmaceutical formulation. When the individual compounds of the combination are administered separately, they are generally each presented as a pharmaceutical formulation as described previously in the art.

Pharmaceutical formulations are often prescribed to the patient in "patient packs" containing the whole course of treatment in a single package. Patient packs have an advantage over traditional prescriptions, where a pharmacist divides a patient's supply of a pharmaceutical from a bulk supply, in that the patient always has access to the package insert contained in the patient pack, normally missing in traditional prescriptions. The inclusion of a package insert has been shown to improve patient compliance with the physician's instructions and, therefore, lead generally to more successful treatment. It will be understood that the administration of the combination of the invention by means of a single patient pack, or patient packs of each component compound, and containing a package insert instructing the patient to the correct use of the invention is a desirable additional feature of the invention.

Hereinafter, the term "active ingredients" means (R,R)-formoterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, preferably (R,R)-formoterol fumarate, and budesonide, or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof.

20

25

30

35

Suitably, the pharmaceutical formulations which are suitable for inhalation according to the invention comprise the active ingredients in amounts such that each actuation provides therapeutically effective dose, for example, a dose of (R,R)-formoterol of 10mcg to 150mcg, preferably 24mcg and a dose of budesonide of 100mcg to 1.6mg, preferably 200mcg to 1mg, more preferably, 200mcg to 400mcg.

The pharmaceutical formulations according to the invention may further include other therapeutic agents for example anti-inflammatory agents such as other corticosteroids (e.g. fluticasone propionate, beclomethasone dipropionate, mometasone furoate, or triamcinolone acetonide), or NSAIDs (e.g. sodium cromoglycate, nedocromil sodium, PDE-4 inhibitors, leukotriene antagonists,

6

iNOS inhibitors, tryptase and elastase inhibitors, beta-2 integrin antagonists and adenosine 2a agonists), or other  $\beta_2$ -adrenoreceptor agonists (such as salbutamol, salmeterol, fenoterol or terbutaline and salts thereof), or anticholinergic agents (such as ipratropium, or tiotropium).

5

10

15

The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous and intraarticular), intranasal, inhalation (including fine particle dusts or mists which may be generated by means of various types of metered dose pressurised aerosols, nebulisers or insufflators), rectal and topical (including dermal, buccal, sublingual and intraocular) administration although the most suitable route may depend upon for example the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredients into association with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

20

. 25

30

Formulations for inhalation include powder compositions which will preferably contain lactose, and spray compositions which may be formulated, for example, as aqueous solutions or suspensions or as aerosols delivered from pressurised packs, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, 1,1,1,2-tetrafluoroethane, carbon dioxide or other suitable gas. Suitable aerosol formulations include those described in EP 0372777 and WO93/11743. For suspension aerosols, the active ingredients should be micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the aerosol formulation, thus the active ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns, for example, 1 to 5 microns.

7

Intranasal sprays may be formulated with aqueous or non-aqueous vehicles with the addition of agents such as thickening agents, buffer salts or acid or alkali to adjust the pH, isotonicity adjusting agents or anti-oxidants.

Capsules and cartridges or for example gelatin, or blisters of for example laminated aluminium foil, for use in an inhaler or insuflator may be formulated containing a powder mix of the active ingredients and a suitable powder base such as lactose or starch. In this aspect, the active ingredients are suitably micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the dry powder formulation, thus the active ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns.

Solutions for inhalation by nebulation may be formulated with an aqueous vehicle with the addition of agents such as acid or alkali, buffer salts, isotonicity adjusting agents or antimicrobials. They may be sterilised by filtration or heating in an autoclave, or presented as a non-sterile product.

Preferred unit dosage formulations are those containing a pharmaceutically effective dose, as hereinbefore recited, or an appropriate fraction thereof, of the active ingredient. Thus, in the case of formulations designed for delivery by metered dose pressurised aerosols, one actuation of the aerosol may deliver half of the therapeutically effective amount such that two actuations are necessary to deliver the therapeutically effective dose.

It should be understood that in addition to the ingredients particularly mentioned above, the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question. Furthermore, the claimed formulations include bioequivalents as defined by the US Food and Drugs Agency.

For a better understanding of the invention, the following Examples are given by way of illustration.

30

15

20

8

## **EXAMPLES**

#### A: Metered Dose Inhalers

## Example 1

5

	Per actuation
(R,R)-formoterol fumarate	24 microgram
Budesonide .	200 microgram
1,1,1,2-Tetrafluoroethane	to 75.0mg

The micronised active ingredients are weighed into an aluminium can, 1,1,1,2-tetrafluoroethane is then added from a vacuum flask and a metering valve is crimped into place.

10

Similar methods may be used for the formulation of Example 2:

## Example 2

	Per actuation	
(R,R)-formoterol fumarate	12 microgram	
Budesonide	100 microgram	
1,1,1,2-Tetrafluoroethane	to 75.0mg	

15

#### **B: Dry Powder Inhalers**

## Example 3

	Per cartridge or blister	
(R,R)-formoterol fumarate	24 microgram	
Budesonide	200 microgram	
Lactose Ph. Eur.	to 12.5mg	
	or to 25.0mg	

9

The active ingredients are micronised and bulk blended with the lactose in the proportions given above. The blend is filled into hard gelatin capsules or cartridges or in specifically constructed double foil blister packs to be administered by an inhaler such as a Rotahaler, Diskhaler, or Diskus inhaler (each of these being a Trademark of Glaxo Group Limited).

Similar methods may be used for the formulations of Example 4:

#### Example 4

10

	Per cartridge or blister
(R,R)-formoterol fumarate	12 microgram
Budesonide	100 microgram
Lactose Ph. Eur.	to 12.5mg
·	or to 25.0mg

#### Claims -

A pharmaceutical formulation comprising (R,R)-formoterol or a
 pharmaceutically acceptable salt, solvate, or physiologically functional
 derivative thereof and budesonide or a pharmaceutically acceptable salt,
 solvate, or physiologically functional derivative thereof, and a
 pharmaceutically acceptable carrier or excipient, and optionally one or
 more other therapeutic ingredients.

10

- 2. A pharmaceutical formulation comprising (R,R)-formoterol fumarate and budesonide, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.
- 15 3. A pharmaceutical formulation according to claim 1 or claim 2 which comprises another corticosteroid, another  $\beta_2$ -adrenoreceptor agonist or an anticholinergic agent.
- 4. A pharmaceutical formulation according to claim 3, wherein the other β<sub>2</sub> 20 adrenoreceptor agonist is salbutamol, salmeterol, fenoterol, terbutaline, or a salt thereof.
  - 5. A pharmaceutical formulation according to claim 3 wherein the anticholinergic agent is ipratropium or tiotropium.

- 6. A pharmaceutical formulation according to any of claims 1 to 5 wherein the amount of (R,R)-formoterol per unit dose is from 87 micrograms to about 150 micrograms.
- 7. A pharmaceutical formulation according to any of claims 1 to 6 wherein the amount of budenoside per unit dose is from above 1.3mg to about 1.6mg.
- 8. A pharmaceutical formulation according to any one of claims 1 to 7 which is suitable for administration by inhalation.

- 9. A pharmaceutical formulation according to any one of claims 1 to 7 which is suitable for intranasal administration.
- 5 10. A pharmaceutical formulation consisting of (R,R)-formoterol or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and optionally one or more other therapeutic ingredients, and 1, 1, 1, 2-tetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane or mixtures thereof as propellant.
  - A method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which a selective β<sub>2</sub>-adrenoreceptor agonist and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a pharmaceutical formulation according to any one of claims 1 to 10.
- 12. A method according to claim 11 wherein the clinical condition is a
  20 disease associated with reversible airways obstruction such as asthma,
  chronic obstructive pulmonary disease (COPD), respiratory tract
  infection or upper respiratory tract disease.
- 13. A Rotahaler, Diskus or Diskhaler inhaler containing a formulation according to any of claims 1 to 8.





PCT7GB 01/01628

CLASSIFICATION OF SUBJECT MATTER PC 7 A61K31/575 A61K A61K31/167 A61P11/06 //(A61K31/575,31:167) According to International Patent Classification (IPC) or to both national classification and IPC **B. FIELDS SEARCHED** Minimum documentation searched (classification system followed by classification symbols) IPC 7 A61K A61P Documentation searched other than minimum documentation to the extent that such documents are included in the fights searched Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, MEDLINE, BIOSIS, CHEM ABS Data, EMBASE, PAJ C. DOCUMENTS CONSIDERED TO BE RELEVANT Citation of document, with indication, where appropriate, of the relevant passages Relevant to claim No. X WO 99 64014 A (ASTRA AB ; EKSTROEM TOMMY 1 - 13(SE)) 16 December 1999 (1999-12-16) claims 1-24 X WO 93 11773 A (ASTRA AB) 1-13 24 June 1993 (1993-06-24) cited in the application claims 1-7 X WO 99 15182 A (TROFAST JAN ; ASTRA AB (SE); 1-13 BAUER CARL AXEL (SE)) 1 April 1999 (1999-04-01) claims 1-10 X 1-13 US 6 004 537 A (CAVANAUGH KELLY A ET AL) 21 December 1999 (1999-12-21) claims 1-24 -/--X Further documents are listed in the continuation of box C. Patent family members are listed in annex. Special categories of cited documents: T later document published after the international filing date or priority date and not in conflict with the application but cled to understand the principle or theory underlying the "A" document defining the general state of the art which is not considered to be of particular relevance Invention "E" earlier document but published on or after the international "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such docu-ments, such combination being obvious to a person skilled "O" document referring to an oral disclosure, use, exhibition or document published prior to the international filing date but later than the priority date claimed in the art. \*&\* document member of the same patent family Date of the actual completion of the international search Date of mailing of the international search report 9 August 2001 04/09/2001 Name and mailing address of the ISA Authorized officer European Petent Office, P.B. 5818 Petentiaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nt, Fax: (+31-70) 340-3016 Herrera, S



Inte II Application No PCT/GB 01/01628

Category *	ation) DOCUMENTS CONSIDERED TO BE RELEVANT  Citation of document, with indication, where appropriate, of the relevant passages	Polyvent to claim No.
Calegory 2	Cuation of coordinate, with indication, where appropriate, of the relevant passages	Relevant to ctaim No.
Υ	WO 98 21175 A (SEPRACOR INC) 22 May 1998 (1998-05-22) claims 19-21	1-13
Y	US 5 795 564 A (MORLEY JOHN ET AL) 18 August 1998 (1998-08-18) abstract	1-13
Y	O'CONNOR B J: "COMBINATION THERAPY" PULMONARY PHARMACOLOGY AND THERAPEUTICS, ACADEMIC PRESS, NEW YORK, NY, US, vol. 11, no. 5/6, 1998, pages 397-399, XP000911059 ISSN: 1094-5539 the whole document	1-13
Y	BOWLER S: "LONG ACTING BETA AGONISTS" AUSTRALIAN FAMILY PHYSICIAN, XX, XX, vol. 27, no. 12, December 1998 (1998-12), pages 1115,1117-1118, XP000973076 the whole document	1-13
X	WO 98 15280 A (TROFAST JAN ;ULLMAN ANDERS (SE); ASTRA AB (SE)) 16 April 1998 (1998-04-16) claims 1-16	1-13
A	DE 23 23 215 A (BOFORS AB) 29 November 1973 (1973-11-29) cited in the application	
A	US 3 929 768 A (BRATTSAND RALPH LENNART ET AL) 30 December 1975 (1975-12-30) cited in the application	
	,	
	·	

ormation on patent family members

tı ıl Ap

Inte al Application No PCT7GB 01/01628

Patent document dited in search report	Publication date	Patent family member(s)	Publication date
WO 9964014 A	16-12-1999	AU 4671099 A BR 9911073 A CN 1305380 T EP 1085877 A NO 20006253 A	30-12-1999 20-02-2001 25-07-2001 28-03-2001 12-02-2001
WO 9311773 A	24-06-1993	AU 673660 B AU 3085892 A CA 2123909 A CZ 9401434 A EP 1101493 A EP 1086697 A EP 0613371 A HR 921445 A HU 75156 A JP 7502036 T NO 942116 A NZ 246050 A SG 48301 A SI 9200403 A SK 73394 A US 5674860 A US 5972919 A	21-11-1996 19-07-1993 24-06-1993 15-12-1994 23-05-2001 28-03-2001 07-09-1994 31-12-1994 28-04-1997 02-03-1995 07-06-1994 21-12-1995 17-04-1998 30-06-1993 08-03-1995 07-10-1997 26-10-1999
WO 9915182 A	01-04-1999	AU 9192898 A BR 9812325 A CN 1271287 T EE 200000145 A EP 1014993 A NO 20001401 A PL 339295 A TR 200000726 T ZA 9808516 A	12-04-1999 05-09-2000 25-10-2000 15-02-2001 05-07-2000 17-03-2000 04-12-2000 21-09-2000 19-03-1999
US 6004537 A	21-12-1999	AU 2194900 A WO 0035441 A	03-07-2000 22-06-2000
WO 9821175 A	22-05-1998	AU 722859 B AU 5175598 A EP 0938467 A US 6040344 A	10-08-2000 03-06-1998 01-09-1999 21-03-2000
US 5795564 A	18-08-1998	US 6068833 A	30-05-2000
WO 9815280 A	16-04-1998	AU 715319 B AU 4578297 A BR 9706822 A CA 2239308 A CZ 9801761 A EP 0871450 A HU 9901674 A JP 2000502365 T NO 982414 A PL 327037 A SK 75198 A	20-01-2000 05-05-1998 23-03-1999 16-04-1998 16-09-1998 21-10-1998 28-09-1999 29-02-2000 27-05-1998 09-11-1998 04-11-1998
DE 2323215 A	29-11-1973	SE 378109 B AT 328630 B	18-08-1975 25-03-1976



Inte al Application No

PCT7GB 01/01628

Patent document cited in search report	Publication date		Patent family member(s)	Publication date
DE 2323215 A		AT	436573 A	15-06-1975
		AU	5525373 A	07~11–1974
		BE	799727 A	17-09-1973
•		CA	1002938 A	04-01-1977
		CH	595400 A	15-02-1978
		CS	178129 B	31-08-1977
		CY	1013 A	23-11-1979
		DD	104295 A	05-03-1974
		DK	134783 B	17-01-1977
		ES FI	414673 A 50631 B	01-07-1976
		FR	2185405 A	02-02-1976 04-01-1974
		GB	1429922 A	31-03-1976
		HK	49179 A	27-07-1979
		HÜ	166680 B	28-05-1975
		IL	42155 A	30-06-1977
		ĴP	1033476 C	20-02-1981
		JP	49041378 A	18-04-1974
		JP	55021760 B	12-06-1980
		KE	· 2970 A	20-07-1979
		NL	7306978 A,B,	21-11-1973
		NO	139640 B	08-01-1979
		SU	470954 A	15-05-1975
		US	3929768 A	30-12-1975
		US	3983233 A	28-09-1976
		YU	129173 A,B	28-02-1981
		ZA	7302955 A	24-04-1974
US 3929768 A	30-12-1975	SE	378109 B	18-08-1975
•		AT	328630 B	25-03-1976
		AT	436573 A	15-06-1975
		AU	5525373 A	07-11-1974
· · · · · · · · · · · · · · · · · · ·		BE	799727 A	17-09-1973
		CA	1002938 A	04-01-1977
		CH	595400 A	15-02-1978
		CS	178129 B	31-08-1977
		CY	1013 A	23-11-1979
		DD	104295 A	05-03-1974
		DE DK	2323215 A 134783 B	29-11-1973 17-01-1977
		ES.	134783 B 414673 A	01-07-1976
		FI	50631 B	02-02-1976
•	,	FR	2185405 A	04-01-1974
		GB	1429922 A	31-03-1976
		HK	49179 A	27-07-1979
		HÜ	166680 B	28-05-1975
		ΪĹ	42155 A	30-06-1977
		ĴΡ	1033476 C	20-02-1981
		JP	49041378 A	18-04-1974
		JP	55021760 B	12-06-1980
•		KE	2970 A	20-07-1979
		NL	7306978 A,B,	21-11-1973
		NO	139640 B	08-01-1979
		SU	470954 A	15-05-1975
		აი	4/0334 //	10-00-19/0
		US	3983233 A	28-09-1976